	Centre Number	Candidate Number
Candidate Name		

## **EXAMINATIONS COUNCIL OF ZAMBIA**

**Examination for General Certificate of Education Ordinary Level** 

# Chemistry

5070/2

Paper 2 Theory

### **Tuesday**

1 AUGUST 2017

Candidates answer on the question paper Additional Information: Mathematical tables/Calculators (non-programmable) Graph paper

#### Time 2 hours

#### Instructions to Candidates

Write your name, centre number and candidate number in the spaces at the top of this page and on any separate answer paper used.

There are **twelve** (12) questions in this paper.

#### **Section A**

Answer all questions.

Write your answers in the spaces provided on the question paper.

#### **Section B**

Answer any three questions.

Write your answers in the separate Answer Booklet provided.

At the end of the examination, fasten your Answer Booklets securely to the question paper.

#### Information for Candidates

The number of marks is shown in brackets [ ] at the end of each question or part question.

The Periodic Table is printed on page 11.

Cell phones are not allowed in the examination room.

FOR EXAMINER'S USI	
Section A	
В9	
B10	
B11	
B12	
TOTAL	

## Section A: [50 marks]

# Answer all questions in the spaces provided.


**A1** The table below shows different gases.

Name of gas	Formula
Carbon monoxide	СО
Helium	He
Methane	CH <sub>4</sub>
Nitrogen	N <sub>2</sub>
Oxygen	O <sub>2</sub>
Sulphur dioxide	SO <sub>2</sub>

(a)	Name	e the gas which will diffuse at the	
	(i)	fastest rate	
		***************************************	
	(ii)	slowest rate	
			[2]
(b)	Which	two gases will diffuse at the same rate?	
		and	[1]
(c)	Which	gas would be expected to diffuse twice as fast as	
	(i)	sulphur dioxide?	
			,
	(ii)	oxygen?	
			[2]
(d)		a gas which will diffuse at a faster rate than any of the gases in above.	
			[1]

[Total 6]

A2 The diagram below shows a chromatogram obtained during the analysis of some substances. Study the chromatogram and answer the questions that follow.



 ${\bf A},\,{\bf B}$  and  ${\bf C}$  are standards whereas  ${\bf P},\,{\bf Q}$  and  ${\bf R}$  are unknown substances.

(a)	which of <b>P</b> , <b>Q</b> and <b>R</b> is a mixture/are mixtures?
	[2]
(b)	Which of the standards can be used to make substance <b>Q</b> ?
	[1]
(c)	Which of the unknown substances has a component whose $R_{\rm f}$ value is different from the $R_{\rm f}$ values of any of the standards?
	[1]
(d)	Explain why the level of solvent is kept below the baseline at the start of the chromatographic process.
	[1]

[Total 5]

**A3** 

Two	elements <b>X</b> and <b>Y</b> are represented by the notations given below.	
	16 7	
	8 Y 3	
Atom	ns of <b>X</b> and <b>Y</b> reacted together to form a compound.	
(a)	What changes in electronic structures occur when atoms of <b>X</b> and <b>Y</b> re	act?
	X:	
	Y:	[2]
(b)	What type of bonding occurs between <b>X</b> and <b>Y</b> atoms?	
		[1]
(c)	Use electron-shell diagrams to show the structure of the compound for between <b>X</b> and <b>Y</b> . Use dots (•) to represent the electrons of <b>X</b> and cre	
724	(x) to represent the electrons of Y. (Show all electron shells)	
		] [2]
(d)	Would you expect the melting point of the compound in (c) above to be	
	high or low? Give a reason for your answer.	•
		[2]
	[Tota	l 7]

A4	the tw PbO <sub>2</sub> ,	o react lead (I	ad (IV) oxide, $Pb_3O_4$ was added to $100cm^3$ of nitric acid solution a tants reacted completely with each other forming lead (IV) oxide, I) nitrate and water. The equation of the reaction is given below. $INO_{3(aq)} \rightarrow PbO_{2(s)} + 2Pb(NO_3)_{2(aq)} + 2H_2O_{(l)}$	nd
	13.70	g of the	e Pb₃O₄ was used in the reaction.	
	(a) Calculate the			
		(i)	mass of lead (II) nitrate formed in the reaction.	
			13	٠
		(ii)	molarity of the nitric acid solution.	-
	(b)	State	with reason whether lead (TV) ovide is a basic ovide or not	[6]
	(b)	State	with reason whether lead (IV) oxide is a basic oxide or not.	
			[Tota	[2] [8]

[Turn over

## Page 6 of 12

A5 The diagram shows part of the Periodic Table. Some elements are represented by letters which are not actual symbols of the elements.

			А		is opposite to	В
С					D	E
F	G		Н	I		J
		К				

Give the letter representing the element which

(a)	has the lowest density	
		[1]
(L)		
(b)	is a liquid at room temperature and pressure.	-,-
		[1]
(c)	forms only a divalent cation.	
	***************************************	[1]
(d)	is used to make electricity cables.	
	3	[1]
(e)	combines with oxygen to form an oxide in which one mole of atoms of t element combines with one mole of oxygen molecules.	he
		[1]
<b>(f)</b>	has atoms with the largest relative atomic mass.	
		Γ <b>1</b>


[Total 6]

•		[Tota	16]
			[1]
	(c)	State one other application of electrolysis.	
		tą.	[2]
	(b)	Write equations for the two reactions which occur at the anode.	
			[3]
	(a)	State the formulae of the ions present in dilute sulphuric acid solution.	
		er. This is in order to increase its resistance to corrosion. This was done be olysing dilute sulphuric acid solution using the aluminium pan as the another.	

An aluminium pan was anodized in order to make the aluminium oxide layer on it

A6

A7 A learner used the experimental arrangement below to investigate the effect of concentration on the rate of the reaction between iron and dilute sulphuric acid.



The learner did the experiment three times using different concentrations of sulphuric acid. The same mass of iron powder was used in each case and the experiments were carried out at a temperature of 20°C.

The results are shown in the table below.

No.	Concentration of sulphuric acid in mol/dm <sup>3</sup>	Volume of hydrogen formed in the first minute/cm <sup>3</sup>
1 .	0.5	14
2	1.0	29
3	1.5	44

(a)		the results, what is the effect of concentration on the rate of on between iron and dilute sulphuric acid?	<b>[1]</b>
(b)	•	in why concentration affects the rate of reaction in the way you described in (a).	F1
			[2]
(c)	State	the effect on the rate of the reaction if	
	(i)	lumps of iron were used instead of the powder.	
	(ii)	the reaction was carried out at 30°C instead of 20°C.	
			[2]

[Total 5]

(a)	Describe any <b>two</b> stages in the industrial manufacture of oxygen.	
		[2]
(b)	Describe the chemical test for oxygen gas.	
88a	***************************************	[2]
(c)	Why must the fractionating column be longer than the one used in separation of crude oil into its various fractions?	
		[1]
(d)	State <b>two</b> industrial uses of oxygen.	
		[2]
	[те	otal 7]

Oxygen is manufactured on a large scale by fractional distillation of liquid air.

**A8** 

## Section B (30 marks)

Answer three questions from this section.

Write your answers in the Answer Booklet provided.

- **B9** (a) A solid substance **S** was suspected to be acidic. Describe what test you would carry out in the laboratory to determine whether **S** was acidic or not.
  - **(b)** When an acid reacts with an alkali, a neutralization reaction occurs and heat is released to the surroundings.
    - (i) Write a balanced chemical equation for the reaction of an acid and an alkali of your choice to form salt and water.
    - (ii) Write an ionic equation, with state symbols, for the reaction in **(b)(i)** above.
    - (iii) Describe how you can obtain crystals of salt from the salt solution formed in the reaction in **(b) (i)**. [7]

[Total 10]

- **B10** Methane is a hydrocarbon whose molecular formula is CH<sub>4</sub>. The gas burns completely in air to produce carbon dioxide, water and heat energy.
  - (a) (i) Write a balanced chemical equation for the complete combustion of methane. [1]
    - (ii) Why should methane never be burnt in limited air? [2]
  - **(b) (i)** Methane is used as a fuel in industry. Using the bond energies in the table below, calculate the enthalpy of combustion of methane.[3]

Bond	Value in KJ/mol
C – C	348
C – H	412
C = 0	831
O – H	464
0 = 0	490

- (ii) Calculate the amount of heat that would be produced from the combustion of 1200cm<sup>3</sup> of methane gas measured at r.t.p. [2]
- (c) Draw the energy level diagram for the combustion of methane. [2]


[Total 10]

You are provided with a sample of copper (II) carbonate (green powder) and any other suitable chemicals and apparatus.

Describe with the aid of a labeled diagram how you can obtain copper metal from the copper (II) carbonate. Write down the chemical equations for the two reactions leading to the formation of the copper metal.

[6]

**(b)** The diagram below shows an experiment which a learner carried out in order to purify copper metal.



- (i) Which electrode, A or B is the impure copper?
- (ii) Name a suitable electrolyte C, used in purification of copper.
- (iii) State **one** physical property of copper and give a reason why copper has this property. [4]

[Total 10]

**B12** (a) The following compounds are members of the homologous series of alkanes.

 $CH_4$ ,  $C_2H_6$ ,  $C_3H_8$ ,  $C_4H_{10}$  and  $C_5H_{12}$ 

- (i) State the formulae of the next **two** members in this series.
- (ii) State with reason whether the compound  $C_9H_{18}$  belongs to the family of alkanes or not.
- (iii) Among the five members given state with reason the one with the highest boiling point. [6]
- **(b)** The demand for petrol worldwide increases every year. To meet the demand, large molecules of hydrocarbons are cracked.
  - (i) What do you understand by the term cracking?
  - (ii) State the **two** types of cracking.
  - (iii) Write down the **two** possible products when the compound  $C_{11}H_{24}$  is cracked. [4]

[Total 10]

[Turn over

DATA SHEET
The Periodic Table of the Elements

Group	0.00							100	22									
<b>;—</b>	=				.72	,						=	2	>	iγ	II>	0	
		a					T Hydrogen				8						He Hellum	
Lí 7	e a					•						ΞŒ	52 CO	‡ Z	<del>α</del> Ο	6 т	20 <b>Ne</b>	
Lithlum	Beryllium 4											Boron 5	Carbon 6	Nitrogen 7		Fluorine 9	Neon 10	555 X 12
	Mg						25					27 <b>A</b> (	ສ <b>ເວ</b> ິ່	با ا		<b>5</b> 29.2	\$ <b>\$</b> }	
=	Magnesium 12											13	14 14	15			18 Augun	
g <b>Y</b>	40 Ca	45 Sc	48 <b>TI</b>	V	JS ZS	ss Mn	56 <b>Fe</b>			n)	86 <b>Zn</b>	70 Ga	გ გ მ	As			월 <b>것</b>	ſ
Potassium 19	7			Vanadium 23	Chromium 24	Manganese 25	tron 26		10000	Capper 29	Zinc 30	Galllum 31	Germanlum 32	Arsenic 33			Krypton 36	
88	88		91	83			104			108	112	115	119	122			131	_
&	s		Zr	g	Mo	ř	Ru	몺		Ag	S		Sn	Sb		×	Xe	
Rubidium 37	Strontium 38	Yltrium 39	Zirconlum 40	Niobium 41	Molybdenum 42	Tchnettum 43	Ruthenium 44	Rhodium 45		Silver 47	Cadmlum 48		22	Antimony 51	Tellurium 52	lodine 53	Xenon 54	W-1000
133	137	139		181	184			-	_	197	201	204		209				_
နှင့် (၁	Ва	La		Ta	N T					Au	Hg	1	요 ]	面	Po	Αŧ	Z	
Caestum 55		57 *	72	73	74	75	76	77	78	92	80 80	81 81	82	83	Polonium 84	Astatine 85	Radon 86	
<b>Fr</b> Francium	226 <b>Ra</b> Radium 88	227 Ac Actinium 89 +									**	1						τ
58-71	58-71 Lanthanoid series	eries		140	141	144		150	152	157	159	162	165	167	169	173	175	
+90-103	-90-103 Actinoid series	ries		S G	Pr	PN PN PN PN PN PN PN PN PN PN PN PN PN P	Pm	Sm	Eu	Gd Gddfinlin	Tb	Dy	Ho	ш	E	ΥÞ	ָר בַּ	
					בן מספרת אווורווו	מפכלוווהבווו	The second second	Colligitain	Teleplan.	פמכלוווני	CINTO	Cyacicaldin				THEIDINIT	רחופותנויי	-

	-y
175 <b>Lu</b> Lutedum 71	<b>Lr</b> Lawrenclu m
173 <b>Yb</b> Yitterbium 70	No Nobelium 102
169 <b>Tm</b> Thullum 69	Mendelevium 101
167 <b>Er</b> Erblum 68	Fm Fermium 100
165 <b>Ho</b> Holmium 67	Es Einsteinium 99
162 <b>Dy</b> Dysproslum 66	Cf Californium 98
159 <b>Tb</b> Terbium 65	<b>BK</b> Berkelium 97
157 Gd Gadoliniu	Curlum 96
152 <b>Eu</b> Europlum 63	Am Ameridum 95
150 Smarium 62	Pu Plutonium 94
Pm Promethium 61	Np Neptunium 93
Neodymium 60	238 <b>U</b> Uranium 92
141 Pr Praseodymlum 59	Pa Protactinium 91
Certum Sertum	232 <b>Th</b> Thorium 90

a = relative atomic mass
X = atomic symbol
b = proton (atomic) number

**key** 

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).

 $N_A = 6.0 \times 10^{23}$ /mol; 1F = 96500C.

# gidemy.com